Contents lists available at ScienceDirect

Results in Physics

journal homepage: www.elsevier.com/locate/results-in-physics

Oscillating dipole with fractional quantum source in Aharonov-Bohm electrodynamics

Giovanni Modanese[a,*]

[a] *Free University of Bolzano-Bozen, P.za Università 5, Bolzano 39100, Italy*

arXiv:1703.05114v1 [physics.gen-ph] 5 Mar 2017

ARTICLE INFO

Article history:
Received ...
Received in final form ...
Accepted ...
Available online ...

Communicated by ...

2000 MSC: 78A25, 78A40, 81Q05

Keywords: Generalized Maxwell theory, Fractional Schrödinger equation, Local current conservation

ABSTRACT

We show, in the case of a special dipolar source, that electromagnetic fields in fractional quantum mechanics have an unexpected space dependence: propagating fields may have non-transverse components, and the distinction between near-field zone and wave zone is blurred. We employ an extension of Maxwell theory, Aharonov-Bohm electrodynamics, which is compatible with currents j^ν conserved globally but not locally; we have derived in another work the field equation $\partial_\mu F^{\mu\nu} = j^\nu + i^\nu$, where i^ν is a non-local function of j^ν, called "secondary current". Y. Wei has recently proved that the probability current in fractional quantum mechanics is in general not locally conserved. We compute this current for a Gaussian wave packet with fractional parameter $a = 3/2$ and find that in a suitable limit it can be approximated by our simplified dipolar source. Currents which are not locally conserved may be present also in other quantum systems whose wave functions satisfy non-local equations. The combined electromagnetic effects of such sources and their secondary currents are very interesting both theoretically and for potential applications.

© 2017 Elsevier B. V. All rights reserved.

Aharonov-Bohm electrodynamics [1] is a natural extension of Maxwell theory which allows to couple the electromagnetic field also to a current density j^μ that is not locally conserved, i.e., it is such that $\partial_\mu j^\mu \neq 0$ in some region ($\mu = 0, 1, 2, 3$). A coupling of this kind would of course be inconsistent in the Maxwell theory, since the Maxwell field equations with source are written as $\partial_\mu F^{\mu\nu} = j^\nu$ and the tensor $F^{\mu\nu}$ is antisymmetric. Aharonov-Bohm electrodynamics has only reduced gauge invariance and one additional degree of freedom, namely the scalar field $S = \partial_\mu A^\mu$ (which is a pure gauge mode in Maxwell theory).

If only locally conserved sources are present, the Aharonov-Bohm theory reduces to Maxwell theory. This happens for all classical sources and for quantum sources which obey a wave equation with locally conserved current, like the Schrödinger equation or Ginzburg-Landau equation. In [2] it was shown, after obtaining a covariant formulation of the Aharonov-Bohm theory and its explicit solution for S, that a censorship property holds: the measurable field strength $F^{\mu\nu}$ does not allow to "see" electromagnetically a non-conserved source j^ν, because it satisfies the equation $\partial_\mu F^{\mu\nu} = j^\nu + i^\nu$, where i^ν is a non-local function of j^ν, called "secondary current" (see eq. (3)), and $j^\nu + i^\nu$ is conserved.

Aharonov-Bohm electrodynamics can be applied in principle to systems which exhibit quantum charge anomalies [3] or to superconducting states described by non-local equations [4, 5]. Very recently, however, a new possible direct application has emerged. As shown by Wei [6], the Schrödinger equation of fractional quantum mechanics [7] has in general a current which is not locally conserved. This may allow particle teleportation and represents a problem for the compatibility of fractional quantum mechanics with Maxwell electrodynamics, but not for Aharonov-Bohm electrodynamics, where instead it leads to new interesting physical possibilities.

In order to illustrate these features in a paradigmatic, simplified situation, we analyze here the features of the Aharonov-

*Corresponding author: Tel.: +39-0471-017134; fax: +39-0471-017009;
e-mail: Giovanni.Modanese@unibz.it (Giovanni Modanese)

Bohm electromagnetic field generated by a non-conserved current of the form

$$j_0 = \left[\delta^3(\mathbf{x}-\mathbf{a}) - \delta^3(\mathbf{x}+\mathbf{a})\right] e^{i\omega t}; \qquad \mathbf{j} = 0. \tag{1}$$

This corresponds to a charge which oscillates periodically between the positions $(\mathbf{x}-\mathbf{a})$ and $(\mathbf{x}+\mathbf{a})$, without an intermediate current, therefore with a kind of teleportation, as allowed by fractional quantum mechanics. The source (1) can be seen as the limit of a suitably defined wave packet (see below).

In order to find the electromagnetic field generated by this source one must solve the Aharonov-Bohm equations, which in covariant form are (with $\partial^2 = \partial_\alpha \partial^\alpha = \partial_t^2 - \nabla$)

$$\partial_\mu F^{\mu\nu} = j^\nu + i^\nu \tag{2}$$

$$i^\nu = -\partial^\nu \partial^{-2} (\partial_\alpha j^\alpha) \tag{3}$$

Let us define, with reference to the four-current (1), the quantity $\theta(t) = \partial_\alpha j^\alpha = \partial_0 j^0$. Eq. (3), which gives the secondary current, can be rewritten as $i^\nu = -\partial^\nu u_{ret}$, where u_{ret} is the mathematical analogue of the retarded electric potential in Lorentz gauge of a charge $\theta(t)$, satisfying the equation $\partial^2 u_{ret} = \theta$. This allows us to compute i^ν at any point P (see Fig. 1). Note that the potential u_{ret} will not be equal to the familiar dipole potential of Maxwell theory, because the source (1) is different from a physical oscillating particle.

From eq. (2) and general unicity theorems ([8]; see also [9]), we deduce that the field $F_R^{\mu\nu}$ at any point R is the Maxwell field with source $j^\nu + i^\nu$. This must include all the contributions from i^ν at each point P, retarded according to the distance RP. Note that such contributions will in general not be transversal with respect to the direction $\overrightarrow{OR}$. The distinction between near field and wave zone will also be blurred, since the secondary source i^ν extends far beyond the localized primary source j^ν.

Finally, let us relate our simplified Ansatz (1) to a non-conserved current in fractional quantum mechanics. According to Wei [6], the correct probability continuity equation in fractional quantum mechanics is $\partial_t \rho + \nabla \cdot \mathbf{j}_a = I_a$, where $1 < a \leq 2$ ($a = 2$ corresponds to ordinary quantum mechanics) and the extra source term is $I_a = -iD_a \hbar^{a-1} \left[\nabla\psi^* (-\nabla^2)^{a/2-1} \nabla\psi - c.c.\right]$. As an example, Wei computes I_a for a wave function of the form $\psi = \psi_1 + \psi_2$, where ψ_1 and ψ_2 are two plane waves with different energies. In order to extend this to a more realistic localized wave function, we define a Gaussian wave packet of the form

$$\psi = \sum_{j=1}^{n+1} \psi_j; \qquad \psi_j = e^{-4(k_j - K)^2 n_1^{-2}} e^{i\hbar k_j x} e^{-i\hbar\omega_j t}, \tag{4}$$

where $k_j = K_0 + j$, $K = K_0 + n_1$, $n_1 = 1 + n/2$. The fractional extra source term can then be written as

$$I_a = -iD_a \hbar^{a-1} \sum_{i,j=1}^{n+1} (1-\delta_{ij}) \left[k_i k_j^{a-1} (\psi_i^* \psi_j - c.c.) + k_i^{a-1} k_j (\psi_j^* \psi_i - c.c.)\right]. \tag{5}$$

As an example, we have computed numerically I_a for $a = 3/2$, at $t = 0$. With parameters $n = 20$ (number of wavelets), $K_0 = 10^3$ (main wave number), $\hbar = 10^{-2}$, $D_{3/2} = 5 \cdot 10^{-3}$ (which simply define the length and mass scale) we obtain the function in Fig. 2. By tuning the parameters we can obtain a wave packet



Fig. 1. Electromagnetic field generated by a four-current $j^\nu(t) = (j^0(t), 0)$ which is not locally conserved. The oscillating dipolar source $\theta(t)$ in O generates a secondary four-current i^ν at any point P in space. i^ν is the four-gradient of $-u_{P,ret}$; $u_{P,ret}$ is mathematically equal to the retarded electric potential generated in P by an equivalent charge $\theta(t)$. The observable field $F_R^{\mu\nu}$ at any point R outside the dipole is found by solving the Maxwell equations with four-current $j^\nu + i^\nu$ at that point (but j^ν is zero in R). Therefore $F_R^{\mu\nu}$ receives "twice retarded" secondary contributions from any P and is not generally transverse to OR.



Fig. 2. Fractional extra-source $I_{3/2}$ for the wave packet (3).

with two peaks more localized in space and a small frequency spread; the result will be approximated well by our Ansatz (1). The full computation of $F_R^{\mu\nu}$ requires in any case a numerical solution.

References

[1] Hively, L.M., and G.C. Giakos (2012) Int. J. of Signal and Imaging Systems Engineering 5, 3; and references therein.

[2] Modanese, G. (2016) Generalized Maxwell equations and charge conservation censorship, arXiv: 1609:00238.

[3] Cheng, T.-P., and L.-F. Li *Gauge theory of elementary particle physics*, Clarendon, Oxford, 2000.

[4] Modanese, G., "Covariant formulation of Aharonov-Bohm electrodynamics and its application to coherent tunnelling", in *Unified Field Me-*

chanics II - Preliminary Formulations and Empirical Tests, Ed.s R.L. Amoroso, L.H. Kauffman, P. Rowlands, World Scientific, 2017.
[5] Clarke, J. (1968) J. de Physique Colloques, 29, C2-3.
[6] Wei, Y. (2016), Phys. Rev. E 93, 066103.
[7] Laskin, N. (2002), Phys. Rev. E 66, 056108.
[8] Woodside, D.A. (2009), Am. J. Phys. 77, 438.
[9] Modanese, G. (2007) Class. Quantum Gravity 24, 1899.